

## LA-UR-21-21793

Approved for public release; distribution is unlimited.

Title: Exploring new small system geometries in heavy ion collisions

Author(s): Lonardonì, Diego

Intended for: One-Page Publication Highlights

Issued: 2021-02-23

---

**Disclaimer:**

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.


# Exploring new small system geometries in heavy ion collisions

## Objectives

- We explore various new collision geometries in the context of the publicly available hydrodynamic model SONIC.
- We incorporate full  $A$ -nucleon configurations for  $^4\text{He}$ ,  $^{12}\text{C}$ , and  $^{16}\text{O}$  obtained from quantum Monte Carlo calculations with realistic nuclear potentials, and use a Monte Carlo Glauber calculation for the initial conditions.

## Impact

- New collision geometries are investigated, with particular focus on p+O and O+O, proposed for running at the Large Hadron Collider (LHC), as well as  $^4\text{He}+\text{Au}$ , C+Au, O+Au, and  $^{7,9}\text{Be}+\text{Au}$ , proposed for running at the Relativistic Heavy Ion Collider (RHIC).
- Hydrodynamic calculations with SONIC indicate flow anisotropies that approximately scale with initial eccentricities, and have an additional dependence on the compactness of the initial geometry (the more compact the source, the larger the flow).
- The comparison between the nucleon distributions obtained from the full 16-nucleon configurations in oxygen and from the  $\alpha$ -cluster tetrahedron model shows significant changes in both p+O and O+O events, suggesting that the new experimental data at LHC should easily be able to discriminate the two cases.



Time evolution of a O+O event (top panel) and  $^4\text{He}+\text{Au}$  event (bottom panel) from SONIC. The color scale indicates the local temperature.

## Accomplishments

S. H. Lim, J. Carlson, C. Loizides, D. Lonardon, J. E. Lynn, J. L. Nagle, J. D. Orjuela Koop, and J. Ouellette, [Phys. Rev. C 99, 044904 \(2019\)](#)